



Diffusion of laparoscopic liver resections: are we there yet?

Nikolaos Machairas, Georgios C. Sotiropoulos

Second Department of Propaedeutic Surgery, National and Kapodistrian University of Athens, Medical School, Athens, Greece

Correspondence to: Nikolaos Machairas, MD, PhD. Second Department of Propaedeutic Surgery, General Hospital Laiko, Ag.Thoma 17, Athens 11527, Greece. Email: nmachair@gmail.com.

Comment on: Goutte N, Bendersky N, Barbier L, *et al.* Laparoscopic left lateral sectionectomy: a population-based study. *HPB (Oxford)* 2017;19:118-25.

Received: 16 March 2018; Accepted: 26 March 2018; Published: 18 April 2018.

doi: 10.21037/ls.2018.04.01

View this article at: <http://dx.doi.org/10.21037/ls.2018.04.01>

Laparoscopic liver surgery has undergone a major evolution during the last 20 years (1). Albeit initially confronted with restraint and skepticism from the surgical community, laparoscopic liver resections (LLR) have gained significant ground and are currently acknowledged as safe and efficient procedures in the hands of hepatobiliary surgeons with experience in laparoscopic surgery (2,3). Over the last years, multiple studies and meta-analyses have evaluated the short- and long-term outcomes of LLR in a plethora of benign and malignant lesions (4-8). Moreover, the laparoscopic approach has been shown to equally merit high-risk groups of patients including cirrhotic, geriatric and obese patients (9-13). LLRs have been associated with improved short-term outcomes as well as equivalent long-term outcomes in the cases of hepatocellular carcinoma (HCC) and colorectal liver metastases (CRLM) (7,14). More recently, the results of the first randomized controlled trial (RCT) on LLR versus open liver resection (OLR) specifically for CRLM were published (15). Parenchyma-sparing LLRs for CRLM were associated with significantly reduced hospital stay and complications' rate compared to OLR while the oncologic adequacy of the procedure was not compromised.

Despite the fact that favorable results for LLR have been clearly documented, the question whether the approach is widely adopted still remains (16). Studies have showed that LLRs are increasingly performed in these specialized centers however others have questioned the diffuse adoption of the laparoscopic approach outside high-volume centers (17-20). To that end, Goutte *et al.* sought to evaluate the adoption and short-term outcomes of laparoscopic left lateral sectionectomy (LLS)—a procedure considered a “gold-standard”—compared to the open approach using a French national database (19). According to their results,

over a 6-year period, only approximately 30% of LLS were performed through the laparoscopic approach. Moreover, laparoscopic LLS was independently associated with a reduced hospital stay regardless of the nature of the hepatic lesion as well as a lower transfusion rate for patients with benign or primary hepatic lesions.

As far as the short-term outcomes in patients who underwent laparoscopic LLS are concerned as presented by Goutte *et al.*, they align with previously published studies (21,22). Liu *et al.* in a recently published meta-analysis, assessed the outcomes from 12 studies which comprised 685 patients who underwent laparoscopic or open LLS (22). Their analysis showed that laparoscopic LLS were associated with a significantly reduced hospital stay ($P<0.001$), lower blood transfusion ($P=0.007$) and a lower morbidity rate ($P=0.01$), compared to the open approach. Another later published study by Goh *et al.* consistently showed reduced hospital stay in patients who had undergone laparoscopic LLS compared to open approach (23).

In the question whether specifically laparoscopic LLS and moreover laparoscopic surgery was widely adopted as a standard procedure by surgeons, Goutte *et al.* state that according to their results, adoption of the laparoscopic approach for LLS was low, and did not show overall improving rates (19). The laparoscopic approach for LLS was performed in 28.5% of the patients and was applied in roughly 33% of the institutions, which performed laparoscopic LLS. Moreover, they showed an increase in the use of the laparoscopic LLS only in university hospitals and high-volume centers (>50 annual liver resections). Studies assessing outcomes from several high-volume centers worldwide have shown significant increase in the

number of LLR performed. Kawaguchi *et al.* showed that LLS (in 27 specialized centers) was performed though the laparoscopic approach in 61.8% of patients compared to 38.2% who underwent open LLS (18). Another recent study from 4 specialized hepatobiliary units showed an average annual percentage change of 12.5% *vs.* -4.1% over a 15-year period for laparoscopic and open procedures, respectively (17).

The small number of cases addressing to peripheral institutions could be partially explained by the limited surgical experience in LLR. Additionally, more complex cases are treated in high-volume centers through the laparoscopic approach rather than more “simple” LLRs, as is the case of LLS. As a result, reduced overall percentage of LLS performed throughout a national healthcare network could be explained. Moreover, laparoscopic LLS performed for a segment II and/or III lesion by a non-proficient surgeon in a peripheral hospital may translate into a more limited and adequate resection when performed by an advanced laparoscopic hepatobiliary surgeon therefore decreasing even more the number of laparoscopic LLS performed in specialized centers.

The way forward for the establishment of LLR are RCTs. Recently the results from the OSLO-COMET trial were published (15). Two more ongoing RCTs, namely the ORANGE II PLUS (NCT01441856) and the ORANGE SEGMENTS (NCT03270917) will further provide evidence on the merits of LLR compared to the traditional open approach. The road to wide diffusion of LLR is still long; the number of these procedures when performed in high volume centers has been shown to increase thus that might not the case in lower volume centers (19,20,24). Acquisition of experience in these procedures outside specialized centers should proceed with caution and is rationally delayed (3,25). Unavoidable centralization of cases in specialized centers ought to emphatically promote stepwise education of younger surgeons in this field (24). Updated data that derive from national databases are mandatory with the aim to further elucidate how the efficient adoption of this approach proceeds over time.

Acknowledgments

Funding: None.

Footnote

Provenance and Peer Review: This article was commissioned

by the editorial office, *Laparoscopic Surgery*. The article did not undergo external peer review.

Conflicts of Interest: Both authors have completed the ICMJE uniform disclosure form (available at <http://dx.doi.org/10.21037/ls.2018.04.01>). The authors have no conflicts of interest to declare.

Ethical Statement: The authors are accountable for all aspects of the work in ensuring that questions related to the accuracy or integrity of any part of the work are appropriately investigated and resolved.

Open Access Statement: This is an Open Access article distributed in accordance with the Creative Commons Attribution-NonCommercial-NoDerivs 4.0 International License (CC BY-NC-ND 4.0), which permits the non-commercial replication and distribution of the article with the strict proviso that no changes or edits are made and the original work is properly cited (including links to both the formal publication through the relevant DOI and the license). See: <https://creativecommons.org/licenses/by-nc-nd/4.0/>.

References

1. Cherqui D. Evolution of laparoscopic liver resection. *Br J Surg* 2016;103:1405-7.
2. Buell JF, Cherqui D, Geller DA, et al. The international position on laparoscopic liver surgery: The Louisville Statement, 2008. *Ann Surg* 2009;250:825-30.
3. Abu Hilal M, Aldrighetti L, Dagher I, et al. The Southampton Consensus Guidelines for Laparoscopic Liver Surgery: From Indication to Implementation. *Ann Surg* 2017. [Epub ahead of print].
4. Sotiropoulos GC, Machairas N, Stamopoulos P, et al. Laparoscopic versus open liver resection for hepatocellular carcinoma: initial experience in Greece. *Ann Gastroenterol* 2016;29:521-9.
5. Ciria R, Cherqui D, Geller DA, et al. Comparative Short-term Benefits of Laparoscopic Liver Resection: 9000 Cases and Climbing. *Ann Surg* 2016;263:761-77.
6. Mirnezami R, Mirnezami AH, Chandrakumaran K, et al. Short- and long-term outcomes after laparoscopic and open hepatic resection: systematic review and meta-analysis. *HPB (Oxford)* 2011;13:295-308.
7. Sotiropoulos GC, Prodromidou A, Kostakis ID, et al. Meta-analysis of laparoscopic vs open liver resection for hepatocellular carcinoma. *Updates Surg* 2017;69:291-311.

8. Sotiropoulos GC, Machairas N, Kostakis ID, et al. Early Experience in Starting a Laparoscopic Liver Resection Program in Greece. *JLS* 2017;21. pii: e2016.00110.
9. Sotiropoulos GC, Prodromidou A, Machairas N. Meta-analysis of laparoscopic vs open liver resection for hepatocellular carcinoma: The European experience. *J BUON* 2017;22:1160-71.
10. Cipriani F, Fantini C, Ratti F, et al. Laparoscopic liver resections for hepatocellular carcinoma. Can we extend the surgical indication in cirrhotic patients? *Surg Endosc* 2018;32:617-26.
11. Sotiropoulos GC, Machairas N, Stratigopoulou P, et al. Laparoscopic liver resection for malignancy in high-risk surgical patients according to ASA classification. *J BUON* 2016;21:1398-402.
12. Machairas N, Kostakis ID, Mantas D, et al. Laparoscopic hepatectomy in a morbidly obese patient with liver cirrhosis: A case report. *Mol Clin Oncol* 2017;6:233-4.
13. Sotiropoulos GC, Machairas N, Kostakis ID. Case Report: Laparoscopic hepatectomy in an elderly patient with major comorbidities. *F1000Res* 2017;6:1286.
14. Zhang XL, Liu RF, Zhang D, et al. Laparoscopic versus open liver resection for colorectal liver metastases: A systematic review and meta-analysis of studies with propensity score-based analysis. *Int J Surg* 2017;44:191-203.
15. Fretland AA, Dagenborg VJ, Bjornelv GM, et al. Laparoscopic Versus Open Resection for Colorectal Liver Metastases: The OSLO-COMET Randomized Controlled Trial. *Ann Surg* 2018;267:199-207.
16. Cleary SP. Minimally Invasive Liver Surgery: Has it Achieved the Standard of Care? *Ann Surg Oncol* 2018. [Epub ahead of print].
17. Berardi G, Van Clevens S, Fretland AA, et al. Evolution of Laparoscopic Liver Surgery from Innovation to Implementation to Mastery: Perioperative and Oncologic Outcomes of 2,238 Patients from 4 European Specialized Centers. *J Am Coll Surg* 2017;225:639-49.
18. Kawaguchi Y, Hasegawa K, Wakabayashi G, et al. Survey results on daily practice in open and laparoscopic liver resections from 27 centers participating in the second International Consensus Conference. *J Hepatobiliary Pancreat Sci* 2016;23:283-8.
19. Goutte N, Bendersky N, Barbier L, et al. Laparoscopic left lateral sectionectomy: a population-based study. *HPB (Oxford)* 2017;19:118-25.
20. Farges O, Goutte N, Dokmak S, et al. How surgical technology translates into practice: the model of laparoscopic liver resections performed in France. *Ann Surg* 2014;260:916-21; discussion 921-2.
21. Bell R, Pandanaboyana S, Hanif F, et al. A cost effective analysis of a laparoscopic versus an open left lateral sectionectomy in a liver transplant unit. *HPB (Oxford)* 2015;17:332-6.
22. Liu Z, Ding H, Xiong X, et al. Laparoscopic left lateral hepatic sectionectomy was expected to be the standard for the treatment of left hepatic lobe lesions: A meta-analysis. *Medicine (Baltimore)* 2018;97:e9835.
23. Goh BK, Chan CY, Lee SY, et al. Laparoscopic Liver Resection for Tumors in the Left Lateral Liver Section. *JLS* 2016;20.
24. van der Poel MJ, Huisman F, Busch OR, et al. Stepwise introduction of laparoscopic liver surgery: validation of guideline recommendations. *HPB (Oxford)* 2017;19:894-900.
25. Yan Y, Cai X, Geller DA. Laparoscopic Liver Resection: A Review of Current Status. *J Laparoendosc Adv Surg Tech A* 2017;27:481-6.

doi: 10.21037/ls.2018.04.01

Cite this article as: Machairas N, Sotiropoulos GC. Diffusion of laparoscopic liver resections: are we there yet? *Laparosc Surg* 2018;2:14.